

Exam 3 Chm 205 (Dr Mattson) 28 March 2018

Academic Integrity Pledge: In keeping with Creighton University's ideals and with the Academic Integrity Code, I pledge that this work is my own and that I have neither given nor received inappropriate assistance in preparing it.

Signature: _____

Name: _____

Chemistry Student Number: _____

Instructions: Show all work whenever a calculation box is provided! Write legibly. Include units whenever appropriate. You will receive credit for **how** you worked each problem as well as for the correct answer. If you need more space, you may use the back of the data sheet provided — Write: "See data sheet" in the answer box — then write your name on the data sheet. On your desk you are allowed only pencils (but no pencil pouch), an eraser, and a non-programmable calculator without a slipcover. Backpacks, bags, and purse-like items must be stored on the tables in the back of the room. Cell phones must be silent and placed in your backpack/bag/purse — not in your pocket.

1. (4 pts) Write the **net ionic** neutralization reaction for a strong acid with a strong base. Include appropriate arrows and a **numerical value** for K_n .

2. (4 pts) Write the **net ionic** neutralization reaction for a weak acid, HA ($K_a = 7.0 \times 10^{-6}$), with a strong base. Include long/short arrows and a **numerical value** for K_n .

3. (4 pts) Write the **net ionic** neutralization reaction for a weak base, A^- , with a strong acid. Include long/short arrows and **numerical value** for K_n ($K_a^{HA} = 6.4 \times 10^{-5}$)

4. (5 pts) Which of these are buffers and if so, what recipe produced the buffer: Circle: **Recipe 1 (R1): wa + wb mixture**, **R2: wa + OH⁻**, **R3: wb + H₃O⁺** or **Not** a buffer.

5.5 mmol KF + 3.0 mmol HNO₃ **R1 R2 R3 Not**

7.0 mmol HF + 7.0 mmol NaOH **R1 R2 R3 Not**

40 mL 1 M KCN + 15 mL 1 M HBr **R1 R2 R3 Not**

5.0 g NaC₂H₃O₂ + 3.0 g HC₂H₃O₂ **R1 R2 R3 Not**

2.0 mmol HClO₂ + 1.2 mmol KOH **R1 R2 R3 Not**


- 5a. (5 pts) What is the pH of a buffer prepared by combining 1.290 g NaNO₂ (MM = 69.00 g/mol) with 500.0 mL 0.100 M HNO₂(aq)? $K_a^{HNO_2} = 3.7 \times 10^{-4}$

Answer with correct significant figures: _____

- 5b. (4 pts) What is the pH of the buffer from Question 5a if 0.0030 mol NaOH is added?

Answer with correct significant figures: _____

6. The following titration curve results when 20.00 mL of an acid is titrated with 0.0994 M NaOH. It takes 33.08 mL to reach the equivalence point.



- 6a. (4 pts) What was the original concentration of the acid? This is a calculation — do not estimate from graph.

Answer with units and significant figures: _____

- 6b. (4 pts) What is K_a for the acid?

Answer with correct significant figures: _____

- 6c. (5 pts) What was the pH after 13.00 mL NaOH has been added? This is a calculation — do not estimate.

Answer: _____

- 6d. (2 pts) From the information provided above, directly or indirectly, including from the graph, do you have everything needed to calculate the pH at the equivalence point? **Circle: Yes or No**

7. (5 pts) Suppose 25.00 mL of a 0.160 M HNO₃(aq) is titrated with 0.205 M KOH(aq). Calculate the pH after 18.00 mL KOH has been added.

Answer to the hundredths place: _____

8a. (4 pt) Calcium fluoride, CaF_2 is a sparingly soluble salt. Write its equilibrium expression and define its K_{sp} in terms the ions produced.

Answer with units: _____

8a. (5 pt) What is the molar solubility of calcium fluoride if $K_{\text{sp}} = 3.5 \times 10^{-11}$? Hint: Set up MICE table.

Answer with units: _____

8b. (4 pts) What is the molar solubility of calcium fluoride in a solution that is 0.040 M KF?

Answer with units: _____

8c. (4 pts) Will a precipitate form if equal volumes of 0.0040 M $\text{Ca}(\text{NO}_3)_2$ and NaF are mixed? Show Q_{sp} calculation! Note: The two solutions dilute each other.

Caution! The solutions dilute each other!

Answer $Q_{\text{sp}} =$ _____ Precipitate: **Yes or No**

Useful equations:

$$R = 8.314 \text{ J/mol K}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$\Delta G = \Delta H - T\Delta S$$

$$\Delta G = \Delta G^\circ + RT \ln Q$$

$$\Delta G^\circ = -RT \ln K$$

9. (5 pts) Which of these processes is entropy-favored?


- dissolving sugar in hot coffee
- mixing of nitrogen gas and bromine vapor
- ethanol boiling
- $\text{N}_2(\text{g}) + 2 \text{H}_2(\text{g}) \rightarrow \text{N}_2\text{H}_4(\text{g})$
- $\text{HCl}(\text{aq}) + \text{Zn}(\text{s}) \rightarrow \text{ZnCl}_2(\text{aq}) + \text{H}_2(\text{g})$

10. (5 pts) Which of these processes are spontaneous?

- dissolving sugar in hot coffee
- mixing of nitrogen gas and bromine vapor
- ethanol boiling at 25 °C and 1 atm
- $\text{CH}_4(\text{g}) + 2 \text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g}) + 2 \text{H}_2\text{O}(\text{g})$
- $\text{HCl}(\text{aq}) + \text{NaOH}(\text{aq}) \rightarrow \text{H}_2\text{O}(\text{aq}) + \text{NaCl}(\text{aq})$

11a. (5 pts) Calculate ΔS° for the reaction in the box, given these standard entropies. Show units in your calculation.

	S° (J mol ⁻¹ K ⁻¹)
$\text{H}_2\text{O}_2(\text{aq})$	144
$\text{H}_2\text{O}(\text{l})$	70
$\text{O}_2(\text{g})$:	205



Answer with units: _____

11b. (4 pts) Given $\Delta H_f^\circ = -188 \text{ kJ}$ for the reaction above, calculate ΔG° for the reaction.

Answer with units: _____

11c. (1 pt) This reaction is spontaneous at...

- all temperatures
- never
- high temperatures only
- low temperatures only


12a. (4 pts) Estimate the boiling point of benzene, given: $\Delta H_{\text{vap}} = 30.7 \text{ kJ/mol}$ and $\Delta S_{\text{vap}} = 87.0 \text{ J mol}^{-1} \text{ K}^{-1}$.

Answer with units: _____

12b. (4 pts) Estimate ΔG_{vap} at 85 °C for benzene. Is vaporization spontaneous at 85 °C?

Answer with units: _____
Is vaporization spontaneous at this temperature? **Yes or No**

13a-d (4 pts). Consider the all-gas reaction at 25 °C.



13a. Is the reaction enthalpy-favored? **Yes or No**

13b. Is the reaction spontaneous at 25 °C? **Yes or No**

13c. Is the reaction entropy-favored? **Yes or No**

13d. Is the reaction ever spontaneous? Use the same answer choices as in Question 11c: **Circle: A B C D**

13e. (4 pts) Calculate ΔG at 25 °C starting with 1.0 atm $\text{P}_4(\text{g})$, 1.0 atm $\text{H}_2(\text{g})$ and 5.0 atm $\text{PH}_3(\text{g})$.

Answer with units: _____


13f. (4 pts) Calculate K_p for the reaction.

Answer: _____

Total score (out of 100): _____

A+ ≥ 95% A ≥ 90% B+ ≥ 85% B ≥ 80% C+ ≥ 75% C ≥ 70% D ≥ 60%

Answers



4. R3, Not, R3, R1, R2

5a. 3.00

5b. 3.09


6a. 0.164 M

6b. $K_a = 1.82 \times 10^{-6}$

6c. 5.55

6d. **Yes**

7. 2.14



8b. $x = 2.06 \times 10^{-4}$ M

8c. $x = 2.19 \times 10^{-8}$ M

8d. $Q_{sp} = 8.0 \times 10^{-9}$ YES

9. a, b, c, e

10. a, b, d, e

11a. $\Delta S^\circ = 57$ J/deg


11b. $\Delta G^\circ = -205$ kJ

11c. A

12a. 353 K

12b. $\Delta G_{\text{vap}} = -0.446$ kJ (kJ/mol is also ok this time)

13a-d (4 pts). Consider the all-gas reaction at 25 °C.



13a. **Yes**

13b. **No**

13c. **No**

13d. **D**

13e. $\Delta G = 45.4$ kJ

13f. $K_p = 6.74 \times 10^{-6}$