

## Exam 2 Chm 205 (Dr Mattson) 27 February 2019

**Academic Integrity Pledge:** In keeping with Creighton University's ideals and with the Academic Integrity Code, I pledge that this work is my own and that I have neither given nor received inappropriate assistance in preparing it.


Signature: \_\_\_\_\_

Name: \_\_\_\_\_

Chemistry Student Number: \_\_\_\_\_

**Instructions:** Show all work whenever a calculation box is provided! Write legibly. Include units whenever appropriate. You will receive credit for **how** you worked each problem as well as for the correct answer. If you need more space, you may use the back of the data sheet provided — Write: "See data sheet" in the answer box — then write your name on the data sheet. On your desk you are allowed only pencils (but no pencil pouch), an eraser, and a non-programmable calculator without a slipcover. Backpacks, bags, and purse-like items must be stored on the tables in the back of the room. Cell phones must be silent and placed in your backpack/bag/purse — not in your pocket.

1. Consider this chart of A going to B. We will build a MICE table from this chart. **Carefully check your work!**




1a. (3 pts) What are the initial concentrations of A,  $[A]_I$ , the change in the concentration of A,  $[A]_C$  and the equilibrium concentration of A,  $[A]_E$ ?

$[A]_I =$	$[A]_C =$	$[A]_E =$
-----------	-----------	-----------

1b. (3 pts) What are  $[B]_I$ ,  $[B]_C$ ,  $[B]_E$ ?

$[B]_I =$	$[B]_C =$	$[B]_E =$
-----------	-----------	-----------

1c. (3 pts) Based on the changes in concentrations of A and B, balance the equation for  $A \rightarrow B$ .



1d. (3 pts) Use the above information to carefully fill in this MICE table. **Be sure you use the balanced equation.**

M	$\rightarrow$
I	
C	
E	

1e. (3 pts) Write the equilibrium constant in terms of  $[A]$  and  $[B]$  **and** then solve for its numerical value.

--

1f. (1 pt) How many seconds does it take for the reaction to reach equilibrium?

1g. (1 pt) If more **A** was added to an equilibrium mixture, how would the reaction shift in order to return to equilibrium? Circle: **Shift left** **Shift right** **No Shift**

1h. (1 pt) Suppose the volume were decreased. How would the reaction shift to return to equilibrium? Circle your choice: **Shift left** **Shift right** **No Shift**

1i. (2 pts) The reaction is **endothermic** and the temperature is increased. What shift occurs to return equilibrium? Circle: **Shift left** **Shift right** **No Shift**

1j. (2 pts) The reaction is **endothermic** and the temperature is increased. How is  $K_C$  affected? Circle:  **$K_C$  increases**  **$K_C$  decreases**  **$K_C$  does not change**

1k. (1 pt) If a catalyst was used and the reaction repeated, how would the time to reach equilibrium change? Circle: **Increase** **Decrease** **No change**

1l. (2 pts) If a catalyst was used and the reaction repeated, how would  $K_C$  change? Circle your choice: **Increase** **Decrease** **No change**

1m. (1 pt) Is  $K_C = K_p$  for this reaction? Circle: **Yes** **No**

1n. (2 pts) At  $t = 4$  s, how does  $Q_C$  compare to  $K_C$ ? Circle:  **$Q_C > K_C$**   **$Q_C < K_C$**   **$Q_C = K_C$**

1o. (3 pts) Write the equilibrium expression for the reverse reaction from Question 1e and solve its numerical value.

--

2a. (4 pts) Consider this equilibrium at 1000 K. Suppose  $[O_2]_E = 0.115$  M. What are  $[SO_2]_E$  and  $[SO_3]_E$ ?

M	2 $SO_2(g)$	+ $O_2(g)$	$\rightleftharpoons$	2 $SO_3(g)$	$\Delta H_{rxn} = -99$ kJ
I	0.200	0.200		0	
C					
E		0.115			

2b. (4 pts) Write the  $K_C$  expression in terms of concentrations. Calculate  $K_C^{1000}$ .

Show work for credit.

2c. (2 pts) How will increasing the temperature affect  $K_C$ ? Circle: **Increase** **Decrease** **No change**


2c. (1 pt) Would decreasing the volume increase  $K_C$ ? Circle: **Yes** **No**

2d. (1 pt) How would decreasing the volume shift the reaction? Circle: **Shift left** **Shift right** **No shift**

2e. (1 pt) Would adding some  $SO_2(g)$  to an equilibrium mixture increase  $K_C$ ? Circle: **Yes** **No**

2f. (1 pt) Would adding  $SO_2(g)$  to an equilibrium mixture cause the reaction to shift right? Circle: **Yes** **No**

3. Consider the reaction profile for the  $A \rightleftharpoons B$  equilibrium. The fraction of molecules,  $f$ , with enough energy to overcome  $E_{act}$  is given



by  $f = e^{-E_{act}/RT}$ . Note: Both  $x$  and  $y$  are positive.

3a. (1 pt) The reaction is **exothermic** or **endothermic**

3b. (1 pt)  $E_{act}^{fwd}$  is designated by... **Circle x or y**

3c. (2 pts)  $\Delta H_{rxn}$  is equal to... **Circle x - y or y - x**

3d. (1 pt) Increasing temperature will result in a ... **Fill in your choice:**  smaller  $E_{act}$  or  larger  $f$  values for both forward and reverse reactions or  change in  $\Delta H_{rxn}$

3e. (1 pt) Increasing temperature will increase  $f$  for... **Fill in your choice:**  the forward reaction more than the reverse reaction or  the reverse reaction more than the forward reaction or  both by the same factor

3f. (2 pts) Increasing temperature will...  increase  $[A]$  or  increase  $[B]$  or  not change  $[A]$  or  $[B]$

3g. (2 pts)  $K_c$  is most likely...   $<1$  or   $>1$  or   $=1$

4. (3 pts) Write the equilibrium expression for water into hydronium ions and hydroxide ions. Use appropriate equilibrium arrows (long/short) given that  $K_w = 1 \times 10^{-14}$  at 298 K. **Include charges on ions for credit on this and all of the following questions.**

5. (3 pts) Select the **three** strong acids from these choices.

- $HNO_2(aq)$      $H_2CO_3(aq)$      $H_2SO_3(aq)$   
  $HClO_4(aq)$      $KNO_3(aq)$      $HCl(aq)$   
  $NH_3(aq)$      $NaBr(aq)$      $HI(aq)$

6. (3 pts) What is the pH of a  $5.8 \times 10^{-4}$  M solution of any monoprotic strong acid?

Answer with correct significant figures: \_\_\_\_\_

7. (4 pts) Write the conjugate base for each weak acid.

a. $HC_2H_3O_2$	b. $HF$	c. $HSO_3^-$	d. $H_3PO_4$
-----------------	---------	--------------	--------------

8. (4 pts) Write the conjugate acid for each weak base.

a. $NO_2^-$	b. $NH_3$	c. $HCO_3^-$	d. $HPO_4^{2-}$
-------------	-----------	--------------	-----------------

9a. (3 pts) Write the equilibrium expression with long/short arrows for propionic acid,  $HC_3H_5O_2$ , a weak acid that you may abbreviate HPr.

9b. (4 pts) What is the pH of a 0.400 M propionic acid solution given that its  $K_a$  value is  $1.35 \times 10^{-5}$ .

Show work for credit.

Answer with correct significant figures: \_\_\_\_\_

10. (3 pts) What is the pH of a  $2.25 \times 10^{-3}$  M  $KOH(aq)$ ?

Answer with correct significant figures: \_\_\_\_\_

11a. (3 pts) A 0.0550 M solution of a weak acid has a pH of 3.94. What is its  $K_a$ ?

Answer with correct significant figures: \_\_\_\_\_

11b. (2 pts) Convert your  $K_a$  value from the previous question into a  $pK_a$  value.

Answer with correct significant figures: \_\_\_\_\_

12a. (4 pts) What is the  $K_b$  value for the acetate ion given the  $K_a = 1.82 \times 10^{-5}$  for acetic acid.

12b. (3 pts) Write the equilibrium expression with long/short arrows for the acetate ion,  $C_2H_3O_2^-$  that you may abbreviate  $Ac^-$ .

12c. (4 pts) Calculate the pH of a  $5.7 \times 10^{-2}$  M  $NaC_2H_3O_2$  solution. (Do not include  $Na^+$  in your MICE table.)

Show work for credit.

Answer with correct significant figures: \_\_\_\_\_


13. (5 pts) Which if these salts are acidic, basic, or neutral? Circle your choice.

- |               |               |              |                |
|---------------|---------------|--------------|----------------|
| A. $NH_4Cl$   | <b>Acidic</b> | <b>Basic</b> | <b>Neutral</b> |
| B. $KI$       | <b>Acidic</b> | <b>Basic</b> | <b>Neutral</b> |
| C. $LiBr$     | <b>Acidic</b> | <b>Basic</b> | <b>Neutral</b> |
| D. $NaBrO_2$  | <b>Acidic</b> | <b>Basic</b> | <b>Neutral</b> |
| E. $Na_2SO_3$ | <b>Acidic</b> | <b>Basic</b> | <b>Neutral</b> |

**Answers:**

1a.  $[A]_I = 0.60 \text{ M}$ ;  $[A]_C = 0.10 \text{ M}$ ;  $[A]_E = 0.50 \text{ M}$

1b.  $[B]_I = 0.0 \text{ M}$ ;  $[B]_C = 0.20 \text{ M}$ ;  $[B]_E = 0.20 \text{ M}$



1d.

M	A	$\rightarrow$	2 B
I	0.60		0.0
C	-0.10		+0.20
E	0.50		0.20

1e.  $K_c = [B]^2 / [A] = 0.080$

1f.  $\sim 15 \text{ s}$

1g. Right

1h. Left

1i. Right

1j.  $K_c$  increases

1k. Decrease

1l. No change

1m. No

1n.  $Q_c < K_c$

1o.  $K_c = [A] / [B]^2 = 12.5$

2a.  $[\text{SO}_2]_E = 0.030 \text{ M}$  and  $[\text{SO}_3]_E = 0.170 \text{ M}$

2b.  $K_c^{1000} = 279$

2c. Decrease

2c. No

2d. Shift right


2e. No

2f. Yes

3a. exothermic

3b. x

3c. x - y

3d.  larger f values for both forward and reverse reactions3e.  the reverse reaction more than the forward reaction3f.  increase [A]3g.   $> 1$ 5.   $\text{HClO}_4(\text{aq})$    $\text{HCl}(\text{aq})$    $\text{HI}(\text{aq})$ 


6. 3.24

7.

a. $\text{C}_2\text{H}_3\text{O}_2^-$	b. $\text{F}^-$	c. $\text{SO}_3^{2-}$	d. $\text{H}_2\text{PO}_4^-$
---------------------------------------	-----------------	-----------------------	------------------------------

8.

a. $\text{HNO}_2$	b. $\text{NH}_4^+$	c. $\text{H}_2\text{CO}_3$	d. $\text{H}_2\text{PO}_4^-$
-------------------	--------------------	----------------------------	------------------------------




9b. 2.63

10. 11.35

11a.  $K_a = 2.4 \times 10^{-7}$

11b. 6.62

12a.  $K_b = 5.5 \times 10^{-10}$



12c. 8.75

13. Acidic, Neutral, Neutral, Basic, Basic